# DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

## LAKE TROPHIC DATA

## MORPHOMETRIC:

Lake: HARRIS	POND	Lake Area (ha):	1.21
Town:	PITTSBURG	Maximum depth (m):	4.8
County:	Coos	Mean depth (m):	
River Basin:		Volume (m³):	22000
Latitude:		Relative depth:	3.9
Longitude:		Shore configuration:	1.03
Elevation (ft	:): 1830	Areal water load (m/yr):	
Shore length	(m): 400	Flushing rate $(yr^{-1})$ :	5.10
Watershed are	ea (ha): 18.1	P retention coeff.:	0.56
% watershed p	onded: 0.0	Lake type: n	atural

BIOLOGICAL:	1 February 1995	2 August 1994
DOM. PHYTOPLANKTON (% TOTAL) #1	DINOBRYON 75%	MALLOMONAS 95%
#2	ASTERIONELLA 10%	
#3	MALLOMONAS 5%	
PHYTOPLANKTON ABUNDANCE (units/mL)		
CHLOROPHYLL-A (µg/L)		40.69
DOM. ZOOPLANKTON (% TOTAL) #1	KERATELLA 41%	KELLICOTTIA 77%
#2	FILINIA 33%	CILIATE SPP. 6%
#3	POLYARTHRA 24%	
ROTIFERS/LITER	369	275
MICROCRUSTACEA/LITER	9	13
ZOOPLANKTON ABUNDANCE (#/L)	378	314
VASCULAR PLANT ABUNDANCE		Sparse
SECCHI DISK TRANSPARENCY (m)		2.7
BOTTOM DISSOLVED OXYGEN (mg/L)	4.5	0.6
BACTERIA (E. coli, #/100 ml) #1		
#2		
#3		

## SUMMER THERMAL STRATIFICATION:

#### stratified

Depth of thermocline (m): 3.0 Hypolimnion volume (m³): None Anoxic volume (m³): 1000

CHEMICAL: Lake: HARRIS POND Town: PITTSBURG						
	1 February 199		2 1	August 199	1994	
DEPTH (m)	1.0	3.0	1.0		3.5	
pH (units)	6.3	6.4	7.0		6.4	
A.N.C. (Alkalinity)	20.8	21.7	11.4		12.9	
NITRATE NITROGEN	0.09	0.08	< 0.05		0.06	
TOTAL KJELDAHL NITROGEN	0.30	0.30	0.38		0.68	
TOTAL PHOSPHORUS	0.015	0.013	0.029		0.098	
CONDUCTIVITY (µmhos/cm)	46.4	47.2	37.6		41.5	
APPARENT COLOR (cpu)	18	23				
MAGNESIUM			1.62			
CALCIUM			3.5			
SODIUM			0.8			
POTASSIUM			0.22			
CHLORIDE	< 2	< 2	< 2		< 2	
SULFATE	5	5	5		5	
TN : TP	26	29	13		8	
CALCITE SATURATION INDEX			2.7			


All results in mg/L unless indicated otherwise

## TROPHIC CLASSIFICATION: 1994

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
**	3	0	6	9	Meso.

#### COMMENTS:

- 1. This is a somewhat remote pond sampled jointly with the NH Fish and Game Department. A very rough, 4-wheel drive road does access the pond.
- 2. Unusually high chlorophyll and phosphorus levels were present for an undeveloped remote pond. Phosphorus was clearly being released from the sediments in the anoxic bottom waters during the summer (note: phosphorus levels were not high in the winter).
- 3. ANC values were relatively high for NH ponds but were similar to nearby Back Lake.



#### FIELD DATA SHEET

LAKE: HARRIS POND

DATE: 08/02/94

TOWN: PITTSBURG

WEATHER: OVERCAST

DATE: 00/02/34	***************************************	ILK: OVERCIBI	
DEPTH (M)	TEMP (°C)	*DISSOLVED OXYGEN	OXYGEN SATURATION
0.1	22.0	8.6	97 %
0.5	22.0	8.6	97 %
1.0	22.0	8.4	95 %
1.5	21.8	7.8	86 %
2.0	21.2	8.5	94 %
2.5	20.0	7.9	85 %
3.0	17.0	0.3	3 %
3.5	14.5	0.4	4 %
4.0	12.0	0.5	5 %
4.5	11.5	0.6	5 %

SECCHI DISK (m): 2.7 COMMENTS:

BOTTOM DEPTH (m): 4.8


TIME:


830

No dissolved oxygen was present below

the thermocline at 3 meters.

\*Dissolved oxygen values are in mg/L





## AQUATIC PLANT SURVEY TOWN: PITTSBURG DATE: 08/02/94 LAKE: HARRIS POND PLANT NAME ABUNDANCE Key COMMON GENERIC Sparse Sweet gale Myrica gale Y Yellow water lily Sparse Nuphar Freshwater sponge Sparse Spongilla 0 d Dulichium arundinaceum Three-way sedge Sparse V Vallisneria americana Tape grass Sparse OVERALL ABUNDANCE: Sparse GENERAL OBSERVATIONS: Freshwater sponges were attached to fallen trees.